

LEAF CURL COPPER FUNGICIDE SPRAY**ACTIVE CONSTITUENT : 125g/kg COPPER (Cu) PRESENT AS COPPER OXYCHLORIDE****Yates Leaf Curl Copper Fungicide Spray works by preventing the entry of the fungus into the plant.****HOW TO USE : DO NOT** apply when soil is dry and plants are suffering from moisture stress.**DO NOT** apply when conditions are hot.**DO NOT** apply if rain is expected within 24 hours.**DO NOT** use on very young seedlings.

PLANT	DISEASE	STATE	RATE PER 1L WATER	HOW TO APPLY
Apples & pears	Black spot (scab)	All states	20g	Apply at green tip.
	Bitter rot	All states except QLD & NT		
Avocados	Anthracoese	NSW, ACT, VIC & QLD only	16g	Apply from flowering to harvest at 28 day intervals.
Beans	Halo blight, common blight	All states	10g	Apply every 10-14 days when disease first appears.
Capsicumis	Bacterial spot	QLD & NT only	16g	Apply every 7-14 days.
		All states except QLD & NT	10g	
Celery	Leaf spot	All states	12-16g	Apply at 7-14 day intervals.
Citrus	Black spot, melanose scab, sooty blotch	All states except QLD, NT & TAS	7g + 5mL white oil	Apply at 50-80% petal fall and again after 6-12 weeks if necessary.
		QLD only	12g + 6.25mL white oil	
	Brown rot	NSW, ACT, VIC & SA only	7g + 5mL white oil	Apply 30-50L spray per tree before onset of autumn rains. If prolonged wet weather period occurs, apply a second spray during the first period of fine weather.
		QLD & NT only	16g	
		WA only	16g + 5mL white oil	
	Brown spot of mandarins	All states except QLD, NT & TAS	8g	Apply at bud burst and 50- 75% petal fall. Further sprays at 6-8 week intervals may be required depending on weather conditions.
Collar rot, Pink disease	QLD & NT only	16g	Apply 30-50L of spray per tree before the onset of Autumn rains.	
Septoria spot	All states except QLD, NT & TAS	7g + 5mL white oil	Apply early to mid March.	
Crucifers (eg. cabbage, cauliflower)	Ring spot, alternaria leaf spot	QLD & NT only	20g	Apply every 14 days.
		All states except QLD & NT	10g	QLD only - do not use on Brussels Sprouts.
Cucurbits (eg. cucumber, melons)	Downy mildew, anthracnose, gummy stem, blight	NSW & ACT only	10g	Apply every 7 days.
Grapes	Downy mildew	All states except QLD	12-16g	Commence when shoots are approx. 20cm long and apply at 14-21 day intervals. Do not apply during flowering.
		QLD & NT only	20g	
Lettuce	Downy mildew, anthracnose	All states	10g	Spray when disease first appears or as a preventative and repeat every 10 days if necessary.
Mangoes	Bacterial black spot	QLD, NT, NSW, ACT, WA only	16g	Apply every 4 weeks from early flowering to fruit set. Ensure thorough coverage.
Ornamentals (Flowers and shrubs)	Black spot (roses only), fungal leaf spots, downy mildew	QLD only	20g	Apply when disease first appears.
		All states except QLD & NT	12g	Repeat at 7-14 day intervals as required.
Passionfruit	Brown spot, septoria spot	QLD & NT only	8g	Apply every 7-14 days.
	Phytophthora blight		12g	
	Brown spot	All states except QLD & NT	16g	
Raspberries	Anthracoese, leaf blight, spur blight	All states	16g	Apply at bud movement.
			10g	Apply just before blossoming, at petal fall and after harvest.
Stonefruit eg. (peaches, nectarines)	Brown rot, leaf curl, freckle, shot hole, rust	All states	30g	Apply at budswell. For shot hole apply again at leaf fall.
Tomatoes and potatoes	Early and late blight (target spot)	All states	20g	Apply at 7-14 day intervals.
Vegetables	Rust, leaf spot diseases	QLD & NT only	20g	Apply at 7-14 day intervals.
		All states except QLD & NT	10g	

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED
DO NOT PICK FRUIT AND VEGETABLES FOR 1 DAY AFTER SPRAYING

HOW TO PREPARE: Add to water and stir thoroughly. The enclosed spoon holds 4 grams when level. Spray evenly over foliage to near run-off, ensuring that both upper and lower leaf surfaces are covered.

CAUTION: Dangerous to fish. DO NOT allow chemical containers or spray to get into drains, sewers, streams or ponds.

STORAGE AND DISPOSAL: Store in the closed, original container in a cool, dry place out of the reach of children. Do not store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.

SAFETY DIRECTIONS: Harmful if swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing the spray wear gloves and a disposable dust mask. Wash hands after use. After each day's use wash gloves.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

yates.com.au



(02) 9763 9200

OR consult



Yates Garden Guide

Yates, a division of Orica Australia Pty Ltd
 1948-1962 Princes Highway, Clayton, VIC 3168, Australia

APVMA No. : 39866 / 0603

BN: DOM: