

**HOW TO USE:** DO NOT mix with other insecticides, fungicides, oils or wetting agents. DO NOT apply in slow drying conditions or if rain is likely before spray is dry. DO NOT apply to wet plants. DO NOT apply when temperatures are expected to be 35<sup>o</sup>C or higher. DO NOT apply when frost is expected. DO NOT apply when soil is dry and plants are suffering from moisture stress.

PLANT	DISEASE	RATE (mL per L water)	HOW TO APPLY
Roses & Ornamentals	Bacterial leaf spot	5mL	Spray at first sign of disease and repeat every 10 – 14 days as required. Do not use on copper-sensitive varieties. To test apply 2 sprays at 14 day intervals.
	Downy mildew	7.5mL	
Beans (including Broad Beans)	Common blight	5mL	Spray at first sign of disease or as a preventative spray. Repeat every 7-14 days as required.
	Bacterial brown spot,		Spray within 21 days of emergence and repeat every 10-14 days depending on disease pressure.
	Chocolate spot, rust		Spray at first sign of disease. Repeat every 10-14 days as required.
	Halo blight		Spray every 10-14 days from when the plants are 15-30cm high, while conditions favour disease.
Beetroot	Downy mildew, rust	5mL	Spray every 10 –14 days from seedling stage until maturity as required.
Brassicas (incl. Broccoli, Brussels Sprouts, Cabbage, Cauliflower)	Downy mildew, black rot, peppery leaf spot, ring spot	5mL	Begin spraying from the seedling stage until maturity. Spray every 10 – 14 days depending on weather conditions and disease pressure. Do not use on copper sensitive varieties.
Capsicums	Bacterial spot, bacterial canker	5mL	Spray at first sign of disease. Repeat every 7-14 days as required. This product will reduce the spread of bacterial canker but will not control seed or soil borne infection.
Carrots	Leaf spot	5mL	Spray at first sign of disease. Repeat every 10-14 days as required.
Celery	Leaf spot	5mL	Spray every 7 – 14 days.
Cucurbits (incl. Cucumbers, Melons, Pumpkin, Zucchini, Squash)	Downy mildew	4mL	Spray when vines are developing or when disease is expected. Only use as a preventative spray. Product will not be effective once the disease is established.
	Angular leaf spot, bacterial leaf spot	5mL	Spray at first sign of disease. Repeat every 7-10 days as required.
Lettuce	Downy mildew, bacterial leaf spot, anthracnose	3mL	Spray at first sign of disease. Repeat every 7-10 days as required. This product predisposes lettuce to frost damage. Lettuce should not be treated with this product if frosts are likely. Do not use on copper sensitive varieties.
Onions	Downy mildew	5mL	Spray at first sign of disease. Repeat every 10 days as required.
Parsnips	Leaf spot	5mL	Spray at first sign of disease. Repeat

			every 10 days as required.
Peas	Ascophyta blight, bacterial blight	5mL	Spray at first sign of disease. Repeat every 10 – 14 days as required.
Potatoes	Early blight / target spot	5mL	Spray from crop emergence to maturity every 7 – 10 days as required. Ensure that both upper and lower leaf surfaces are sprayed.
Rhubarb	Crown rot	5mL	Dip rhubarb crowns before planting
	Downy mildew		Spray every 14 days from seedling stage until maturity as required.
Silver beet	Downy mildew	5mL	Spray from seedling stage to maturity every 10-14 days as required. Do not use on copper sensitive varieties.
Spinach	Downy mildew	5mL	Spray at first sign of disease and repeat every 10 – 14 days as required.
Tomatoes	Bacterial spot, bacterial speck, bacterial canker	5mL	Apply as a preventative spray every 7 – 10 days. Ensure thorough leaf coverage and that both upper and lower leaf surfaces are covered. This product will reduce the spread of bacterial canker but will not control seed or soil borne infection.
	Early blight / target spot		Begin spraying shortly after transplanting when plants are 15cm high and before symptoms appear. Apply every 7 – 10 days as required. Ensure that both upper and lower leaf surfaces are sprayed.
	Late blight / Irish blight		Spray at first sign of disease and repeat every 10 – 14 days.
Apples	Black spot (scab)	5mL	Spray at green tip before tips are 1cm. Ascertain the tolerance of recently introduced varieties to copper sprays.
Apricots	Shot hole, freckle	5mL	Spray at budswell prior to signs of leaf / bud movement. Also spray at least once after harvest.
	Bacterial gummosis	6.25mL 5mL (NSW only)	Autumn: spray at 25-50% leaf fall and again at 90-100% leaf fall. Winter: spray in mid winter. Spring: spray at first sign of bud movement. Repeat application 7-10 days later.
		3.25mL (NSW, VIC, SA, WA only)	Spray 1 week after petal fall. Repeat 7-10 days later. These sprays control the high leaf population of the bacteria in mid to late spring.
Avocadoes	Anthrachnose, cercospora spot, sooty blotch (QLD, NSW, VIC, SA, WA only)	5mL	Spray every 4 weeks from the end of flowering to harvest. During extended wet weather spray every 14 days.
Cherries	Shot hole	5mL	Spray when buds are opening but before and within one week of bud opening.
	Bacterial gummosis / canker	6.25mL (VIC, TAS, SA, WA only)	Autumn: spray at 25-50% leaf fall. Apply again at 90-100% leaf fall. Winter: spray in mid winter. Spring: spray at first sign of bud

		5mL (NSW only)	movement.
		3.25mL (NSW, VIC, SA, WA only)	Spray 1 week after petal fall. Repeat application 7 – 10 days later. These sprays control the high leaf population of the bacteria in mid to late spring.
Grape vines	Downy mildew, powdery mildew	5mL	Spray when shoots are 10cm long and repeat at 10 to 14 day intervals or as necessary while weather conditions favour infection. Leaf damage may occur on copper sensitive varieties.
Lychees	Lychee pepper spot	5mL	Spray every 4 weeks from the end of flowering to harvest. During extended wet weather, spray every 14 days.
Macadamia	Husk spot (QLD, NSW, WA, NT only)	5mL	Good spray penetration of foliage is essential. Apply from nut set (late Sept) to Dec. Apply at least 3 sprays at 3-4 week intervals.
	Anthrachnose (QLD, NT, NSW only)		Good coverage inside the tree is essential. Spray from early summer to May at monthly intervals.
	Pink limb blight		Good coverage of infected limbs from early summer to May at monthly intervals.
Mangoes	Anthrachnose	7.5mL (NSW, QLD, SA, WA, NT only)	Spray every 4 weeks from end of flowering to harvest. During extended wet weather spray every 14 days. Use in rotation with other fungicides.
	Bacterial black spot	5mL	Spray at 2-4 week intervals from just after fruit set until harvest.
Nectarines, Peaches, , Almonds	Shot hole	5mL	Spray when buds are swelling but before and within one week of bud opening.
	Leaf curl		Correct timing is critical for effective control. Spray when buds are swelling but before and within one week of bud opening. Where leaf curl is likely to be a severe problem based on previous experience, spray first in autumn at leaf fall and then at the first sign of bud swell and repeat one week later.
Pears	Black spot (scab)	5mL	Spray at green tip before tips are 1cm. Ascertain the tolerance of recently introduced varieties to copper sprays.
Plums	Shot hole	5mL	Spray when buds are swelling but before and within one week of bud opening.
Strawberries	Leaf spot, grey mould	5mL	Spray when plants are established and repeat every 7-10 days. Discontinue if reddening of leaf veins occurs or new growth is inhibited.

Avocadoes, Bananas (NSW, WA only), Citrus, Kiwifruit, Lychees, Nectarines, Peaches, Plums, Passionfruit, Tropical fruit, Macadamia (QLD, WA only), Pecans	Phytophthora stem canker	250mL/L water or water based paint	Mix thoroughly. Apply only to stems of trees where cankers appear, after removing dead tissue. Repeat applications up to a maximum of 5 per season until natural healing has commenced. If applied mixed with a water-based paint, one or two treatments per season may be all that's required.
---	--------------------------	------------------------------------	---

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED**

Withholding Period: **DO NOT PICK GRAPES FOR 4 WEEKS AFTER SPRAYING**  
**DO NOT PICK ALL OTHER CROPS FOR 1 DAY AFTER SPRAYING**

**STORAGE & DISPOSAL :** Store in the closed, original container in a cool, dry place out of reach of children. Do not store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.

® Yates is a registered trademark. The front panel of this package is owned under copyright laws by Yates Australia.

™ The Squares device is a trade mark.



**LIABILITY:** This product must be used strictly as directed. Yates will not be liable for any loss or damage arising from a failure to strictly follow the directions for use.